

Mix & Gro

Professional Quality Mulch, Soil Amendment and Potting Mix

People and Pet Safe



Contains Organic Based Ingredients

**Earth Friendly
Natural Based
Formula**

Blue Ribbon Blend

Contains naturally occurring microbial diversity. Nourishes soil-inhabiting organisms, saves water and serves as a source of support for planting and managing soil structure. Made from completely composted mushroom compost substrate. And is free from GMOs, pesticides, growth hormones, sewage sludge and synthetic chemicals. For use with seeds, starter plants, established plants, trees, shrubs, lawns or as a cover mulch.

“Not for use in organic crop and organic food production in the State of California”

What Is Mushroom Compost?

Mushroom compost is the growing material that's left over from the process of growing mushrooms on an industrial scale. It's sometimes referred to as '**spent mushroom compost**'.

Why Is It Useful?

There are plenty of reasons to use mushroom compost in your garden and only a few reasons why you shouldn't. Let's start with the reasons why:

1) It feeds your plants

Mushroom compost is a slow release nourishing fertilizer. It supplies beneficial fungi, bacteria and micro-organisms into the soil. If you spread mushroom compost through the garden or used it as mulch, your plants probably won't need additional fertilizer feedings.

2) It's versatile

You can mix mushroom compost into vegetable beds prior to planting or you can put it around plants that are in situ – this is known as 'top-dressing'.

3) It's organic

Mushroom compost is mostly derived from organic substrates including high quality peat moss. You won't need to worry about chemicals leaching into your veggies or affecting wildlife.

4) It holds water

Mushroom compost is organic matter and holds moisture. We're all trying to cut back on our water usage, so soil that can retain moisture is good for the environment.

Moisture retention also improves the life of your plants because it stabilizes the changes between dry to wet seasonal conditions.

5) No Pests

Mushroom compost is steam pasteurized before its removed from the mushroom growing house. It will not introduce pests or pathogens into your garden.

6) Mushroom compost is guaranteed weed-free.

7) No viruses

Pests and weeds are not the only issues our plants have to fight off. Viruses can kill plants quickly. Mushroom compost is virus-free and it helps to produce a vigorous plant that is more able to can fight off infections.

8) It prevents surface compaction

Soil surface is easily compacted by sun, rain, and foot tread, but mushroom compost can prevent the soil surface from crusting over and compacting. This aids rain absorption and encourages worms to generate healthy and a crumbly soil.

9) It's a drainage aid

Mushroom compost can improve chalky, clay or compacted soils so that water runs freely. This means your plants won't get waterlogged and drowned out. Rotting roots can destroy the toughest of plants.

10) It has super weed suppressant powers

If you apply a thick layer of mushroom compost as mulch it blocks out light and kills off weeds. This is a big bonus for the busy gardeners who don't have time to deal with nutrient stealing dandelions, creeping buttercup and mare's tail.

11) It doesn't smell bad

Yes, mushroom compost is made from organic matter, but it doesn't smell bad. When you first open the bag, it may smell musty or 'dark' but it certainly doesn't stink. Once it's on the garden any lingering smell disappears.

Why You Shouldn't Use Mushroom Compost

There are a few plants that are not keen on the alkaline properties of mushroom compost. This includes fruit bushes and acid loving plants like camellia, rhododendron, heathers and magnolia. If you have put mushroom compost around these plants and noticed yellowing leaves it points to a nutrient deficiency. Dig out the mushroom compost and get some acid-based compost in there promptly.

Plant Nutrients and Fresh Mushroom Compost

Dr. Mike Fidanza, Associate Professor of Biology (Plant and Soil Sciences), The Pennsylvania State University, Berks Campus, Reading, PA, Email: fidanza@psu.edu and

Dr. David Beyer, Professor of Plant Pathology, The Pennsylvania State University, University Park, PA, Email: dmb8@psu.edu

The purpose of this research project was to measure the plant nutrient content and particle size distribution of fresh mushroom compost. Mushroom compost, formerly referred to as “spent mushroom substrate” or “SMS,” is the composted organic material remaining after a mushroom crop is harvested. Although there have been a few scattered reports and observations on the chemical compounds found in mushroom compost that are useful for the growth of agricultural crops and other plants, no formal record exists specifically for fresh mushroom compost. The key word is “fresh” – the material obtained directly as it is removed from a commercial mushroom production facility and not “static-aged” by being stockpiled outdoors in a field for several months.

During late winter/early spring 2005, 30 fresh mushroom compost samples were collected from mushroom farms in Berks and Chester counties. Each sample was placed in a one-gallon plastic container, sealed and sent to the Agricultural Analytical Services Laboratory (Pennsylvania State University, University Park, PA) for processing and analysis. For this study, fresh mushroom compost samples were processed and analyzed, and results are presented on a wet weight basis, wet volume basis, and dry weight basis (Table 1), particle size distribution (Figure 1), and amount of plant nutrients on a per acre basis (Table 2).

pH Most agricultural and horticultural crops grow best within a soil pH range of 6.0 to 7.0 (i.e., < 7.0 is acidic, 7 is neutral, and > 7 is alkaline). Within this pH range, most nutrients in the soil exist in an available form that can be taken-up by plant roots. Keep in mind, there are exceptions. For example, blueberries prefer a more acidic soil pH. The average pH of fresh mushroom compost is 6.6, an excellent pH for any compost used as an organic fertilizer or soil amendment. Unfortunately, rumors have bounced around for years about the pH of mushroom compost being too acidic or too alkaline for growing plants but this is not the case.

Soluble Salts This statement has been repeated many times over the years: “...you can’t use mushroom compost because of the high salt content.” With soils and composts, the salts of concern are those positively charged cations: potassium (K^+), calcium (Ca^{2+}), magnesium (Mg^{2+}) and sodium (Na^+). An excessive amount of these salts dissolved in the soil solution (i.e., the soil water environment) can increase the osmotic pressure of the soil solution, and this “salt effect,” also referred to as salinity, inhibits water absorption by seeds and roots. Many composts and fertilizer products contain these salts in varying amounts. Potassium, calcium, and magnesium are actually essential nutrients beneficial to growing plants. When adding compost

or fertilizer to soil, these salts are often diluted by leaching with adequate rainfall or irrigation, or by tilling or mixing those materials into the soil.

Soluble salt content in soil and compost is measured indirectly by electrical conductivity, and the methods vary with each laboratory. Penn State's laboratory determines soluble salts using a 1:5 (compost:water) slurry. The average soluble salt content of fresh mushroom compost is not in an amount high enough to cause problems with plant growth. With fresh mushroom compost or any other compost or fertilizer, however, over-application or incorrect application of these materials to the soil can result in an excessive salt load.

Excess sodium salt in soil can result in problems with soil structure and drainage as well as inhibiting water absorption by plant roots. The best way to address this issue with fresh mushroom compost or any compost or organic soil amendment is to calculate the sodium adsorption ratio (SAR) of the product or material. The SAR compares the sodium concentration relative to the concentrations of calcium and magnesium. The SAR is calculated as follows:

$$\text{SAR} = \frac{[\text{Na}^+]}{\sqrt{\frac{([\text{Ca}^{2+}] + [\text{Mg}^{2+}])}{2}}}$$

A SAR value ≥ 15 indicates an excess amount of sodium compared to calcium and magnesium, and that sodium would be adsorbed by the soil clay particles thus causing problems mentioned above. Applying 40 tons of fresh mushroom compost to one acre of land (calculated by using a bulk density amount of 575 lbs/yd³) results in a SAR = 0.38, which is very low! Therefore, the presence of sodium in fresh mushroom compost is not a negative aspect of this product, since there is an ample amount of calcium and magnesium present to prevent sodium from accumulating on those soil particles.

The *bottom line* with fresh mushroom compost, or any compost or organic soil amendment or fertilizer, is *environmental stewardship*. Compost products used for agricultural crop production, horticulture plant production, gardening, or land use reclamation should be applied correctly and in the proper amount. For many years, mushroom compost was mislabeled as "mushroom soil," and the product was unfortunately treated like a soil. As a result, Pennsylvania's mushroom industry had to deal with the negative feedback of trying to explain why their mushroom compost was not behaving like topsoil. Mushroom compost is not topsoil, rather an excellent compost useful to improve soil health and plant growth.

Bulk Density, Solids and Moisture The average bulk density of fresh mushroom compost is essentially 575 lbs/yd³ (wet volume basis), with over half of the overall weight attributed to water. Fresh mushroom compost contains solids at 42.7 percent (wet weight) or 243.4 lbs/yd³ (wet volume), and moisture or water at 57.3 percent (wet weight) or 331.5 lbs/yd³ (wet volume). The ideal moisture content of compost depends on the water holding capacity of materials used to produce the compost. Overall, composts higher in organic matter have a higher water holding capacity. A range of 35 to 55 percent (wet weight) for solids and 45 to 65 percent

(wet weight) for moisture is ideal for most compost products. Fresh mushroom compost falls into those ranges.

Organic Matter and Carbon The average organic matter content of fresh mushroom compost is 26 percent (wet weight) or 147 lbs/yd³ (wet volume). Fresh mushroom compost is an excellent source of organic matter, which represents a pool of plant nutrients to be slowly released over time. Also, due to the high organic matter and carbon content, fresh mushroom compost would be extremely useful to amend soils low in organic matter and nutrient availability, especially sand-based soils.

Carbon:Nitrogen (C:N) ratio The amount of carbon relative to the amount of nitrogen is an indicator of nitrogen availability for plant growth. The ideal C:N ratio for good composts should be within the range of 10:1 to 15:1, and no greater than 30:1. At higher C:N ratios, soil microorganisms can immobilize or tie-up nitrogen making it unavailable for plant roots. The average C:N ratio for fresh mushroom compost is ideal at 13:1.

Primary Macronutrients Nitrogen (N), phosphorus (P) and potassium (K) are important and essential primary plant macronutrients needed in higher quantities by plants than other nutrients. The average total N content of fresh mushroom compost is 1.1 percent (wet weight) or 6.4 lbs/yd³ (wet volume). The majority of this N is in the organic form, with a very small percentage in the ammonium-form. In general, all organic compost materials (for example, composts made from landscape and yard wastes, plant residues, animal wastes) have low N content usually in the 1 to 3 percent range. Compost is a natural organic source of N, and the N is released slowly by soil microbial decomposition. Plants use N for growth and development, especially for amino acid and protein synthesis, and also for chlorophyll production. The average phosphate (phosphorus in the form of P₂O₅) content of fresh mushroom compost is 0.7 percent (wet weight) or 3.8 lbs/yd³ (wet volume). Phosphorus is needed in plants for cell energy transfer and electron transport, and for DNA and RNA synthesis. Also, phosphorus is essential for seed germination and emergence. The average potash (potassium in the form of K₂O) content of fresh mushroom compost is 1.3 percent (wet weight) or 7.1 lbs/yd³ (wet volume). Potassium is used by plants for enzyme reactions and the osmotic regulation of cells.

Secondary Macronutrients Calcium (Ca), magnesium (Mg), and sulfur (S) are considered secondary plant macronutrients, and are also required by most plants, but not in large quantities like the primary macronutrients of N, P, or K. Fresh mushroom compost contains Ca at 2.3 percent (wet weight) or 13.2 lbs/ yd³ (wet volume), Mg at 0.4 % (wet weight) or 2.0 lbs/ yd³ (wet volume), and S at 0.9 percent (wet weight) or 4.9 lbs/ yd³ (wet volume). Calcium is important in plants for cell membrane structure and function. In plants, Mg is a central component of chlorophyll and vital for photosynthesis, and S is important for amino acid synthesis.

Micronutrients Iron (Fe), manganese (Mn), copper (Cu), and zinc (Zn) are all considered plant micronutrients and are needed in much smaller quantities compared to the macronutrients. Sodium (Na) and aluminum (Al) are not typically listed as micronutrients but are included in most compost analysis tests. All of these nutrients are available in fresh mushroom compost at a very low average range of 0.01 to 0.2 percent (wet weight) or 0.03 to 1.1 lbs/yd³ (wet volume). Refer to Table 1 for the exact amounts of each nutrient. In plants, chlorophyll synthesis (Fe), formation of oxygen during photosynthesis (Mn), cellular respiration (Cu), and enzyme functions (Zn) are supported by these micronutrients. Again, rumors of excessive or toxic amounts of zinc present in fresh mushroom compost are not accurate as these results indicate.

Particle Size Approximately 91 percent of fresh mushroom compost is $\leq 3/8$ inches in diameter (Figure 1). Therefore, fresh mushroom compost has a consistent and uniform size, which translates to ease of transport and application. Fresh mushroom compost is not “clumpy” or difficult to handle.

So, how much of these plant nutrients are supplied from fresh mushroom compost on a per acre basis? To apply evenly one-inch thick fresh mushroom compost to one acre of land would require 40 tons of fresh mushroom compost as calculated from an average bulk density of 575 lbs/yd³ (Table 2). This calculation shows a total nitrogen amount of 891 lbs, of which 29 lbs is quickly available nitrogen (ammonium-nitrogen) used immediately by a crop in the same growing season when this compost is applied. A remaining amount of 862 lbs of organic nitrogen represents nitrogen that is slowly released over time. A typical “rule of thumb” is that 10 to 20 percent (86 to 192 lbs) of nitrogen could potentially become available during the growing season from this organic nitrogen pool. This kind of information is useful in field crop production in order to calculate nitrogen supplied by compost and nitrogen needed from fertilizer inputs. With the recent increase in synthetic fertilizer costs, nitrogen supplied from fresh mushroom compost represents an economical way to meet crop nutrient needs while minimizing the expense of applying synthetic fertilizers. Phosphate information on a per acre basis is also useful, since some states require detailed nutrient management plans for the purpose of monitoring the amount of phosphate being applied to the land.

In conclusion, fresh mushroom compost applied to soil or incorporated into soil has many benefits: improves soil structure, provides plant nutrients, increases plant nutrient availability, increases soil microbial populations, increases soil cation exchange capacity, increases plant root structure, increases soil aeration, improves soil water status, and reduces soil compaction. Fresh mushroom compost is a viable “green” product as an organic soil amendment and fertilizer for crop production systems and other land management issues.

For more information on the cost of Penn State’s compost analysis and other related information, refer to the laboratory Web site at www.aasl.psu.edu. Also, before sending compost samples to any laboratory, make sure it is U.S. Compost Council certified (www.compostcouncil.org).

Acknowledgements

Funding was provided by the Mushroom Industry Farmer-Based Applied Research program administered through the AMI (Washington, DC) and by the Chester County Economic Development Council, Project #C000029029 (Exton, PA).

CAC's Mushroom Compost Committee under the direction of Tom Brosius, Marlboro Mushrooms; Don Needham, Hy-Tech Mushroom Compost, Inc.; and Eugene D. Richard, Richard Enterprises Inc., provided technical support for this research project.

Committee Members:

Chip Chalupa, Modern Mushroom Co

Phil Coles, Giorgi Mushroom Co.

Joe DiNorscia, Laurel Valley Soils

Linda Farrell, Blue Heron Consulting

Joseph Fecondo, Joseph Silvestri & Sons

Dan Greig, Chester County Conservation District

Charles Krallinger, Giorgi Mushroom Co.

Dr. Kenneth Lomax, University of Delaware

Sara Manning, CAC/AMI

Don Moore, Agronomist, Agricultural, Environmental & Technical Consulting (AET)

Mike Pia, Kaolin Mushrooms

Laura Phelps, American Mushroom Institute

Joseph G. Poppiti, JoBeth Farm

Chris Strohmaier, Chester County Conservation District

David Tranquillo, Giorgi Mushroom Co.

Further Reading

For more information on soils, refer to these publications:

Brady, N.C. and R.R. Weil. 2000. Elements of the nature and properties of soils. Prentice Hall, Upper Saddle River, NJ.

Brady, N.C. and R.R. Weil. 1996. The nature and properties of soils. Prentice Hall, Upper Saddle River, NJ.

Foth, H.D. 1984. Fundamentals of soil science. John Wiley and Sons, New York, NY.

Miller, R.W. and D.T. Gardiner. 2001. Soils in our environment. Prentice Hall, Upper Saddle River, NJ.

Singer, M.J. and D.N. Munns. 2002. Soils, an introduction. Prentice Hall, Upper Saddle River, NJ.

Sidebar: Success Story!

The Pennsylvania Department of Agriculture had classified mushroom compost (formerly listed as “spent mushroom substrate” or “SMS”) as an agricultural waste product, which then involved regulation through the Pennsylvania Department of Environmental Protection. This classification was incorrect, and resulted in unfortunate environmental and economic challenges for Pennsylvania’s mushroom industry. As a result of this research by Drs. Mike Fidanza and David Beyer, and CAC’s Mushroom Compost Committee, mushroom compost has been reclassified correctly as a fertilizer and soil amendment. For a copy of a fertilizer/soil amendment label for fresh mushroom compost, refer to the website www.mushroomcompost.org or AMI’s website www.americanmushroom.org.

Table 1. Average values from analysis of fresh mushroom compost on a wet weight basis, wet volume basis, and dry weight basis.

<u>Parameter Measured⁽¹⁾</u>	<u>Wet Weight Basis⁽²⁾</u>	<u>Wet Volume Basis⁽²⁾</u>	<u>Dry Weight Basis⁽²⁾</u>
pH	6.6	---	---
Soluble Salts ⁽³⁾	13.3 mmhos/cm	---	---
Bulk Density	---	574.7 lbs/yd ³	---
Solids	42.7 %	243.4 lbs/yd ³	---
Moisture	57.3 %	331.5 lbs/yd ³	---
Organic Matter	25.9 %	146.7 lbs/yd ³	61.0 %
Carbon	14.3 %	81.1 lbs/yd ³	33.4 %
Carbon:Nitrogen Ratio	12.8:1 (~13:1)	12.8:1 (~13:1)	12.8:1 (~13:1)
Total Nitrogen	1.1 %	6.4 lbs/yd ³	2.7 %
Organic Nitrogen	1.1 %	6.2 lbs/yd ³	2.6 %
Ammonium Nitrogen (NH ₄ -N)	0.03 %	0.2 lbs/yd ³	0.08 %
Phosphate (P ₂ O ₅)	0.7 %	3.8 lbs/yd ³	1.6 %
Potash (K ₂ O)	1.3 %	7.1 lbs/yd ³	2.9 %
Calcium	2.3 %	13.2 lbs/yd ³	5.4 %
Magnesium	0.4 %	2.0 lbs/yd ³	0.8 %
Sulfur	0.9 %	4.9 lbs/yd ³	2.0 %
Sodium	0.1 %	0.7 lbs/yd ³	0.3 %
Aluminum	0.1 %	0.9 lbs/yd ³	0.3 %
Iron	0.2 %	1.1 lbs/yd ³	0.4 %
Manganese	0.02 %	0.1 lbs/yd ³	0.04 %
Copper	0.01 %	0.03 lbs/yd ³	0.01 %
Zinc	0.01 %	0.05 lbs/yd ³	0.02 %

⁽¹⁾Fresh mushroom compost samples ($n = 30$) collected in one-gallon size amounts were analyzed by the Agricultural Analytical Services Laboratory (Pennsylvania State University, University Park, PA), from January through April 2005.

⁽²⁾Mushroom compost samples analyzed "as is" when received at the laboratory for wet weight and wet volume measurements; for dry weight basis, samples oven-dried to remove moisture, then analyzed.

⁽³⁾Soluble salts determined by measuring electrical conductivity in a 1:5 (compost:water, weight ratio) slurry.




Figure 1. Average diameter values for particle size distribution of fresh mushroom compost as determined from a wet weight basis. Fresh mushroom compost samples ($n = 30$) collected in one-gallon size amounts were analyzed by the Agricultural Analytical Services Laboratory (Pennsylvania State University, University Park, PA), from January through April 2005.

Table 2. Amount of plant nutrients from 40 tons of fresh mushroom compost applied to one acre of land.

<i>Parameter⁽¹⁾</i>	<i>Amount (lbs)⁽²⁾</i>
Solids	33,877
Moisture	46,140
Organic Matter	20,425
Carbon	11,294
Total Nitrogen	891
Organic Nitrogen	862
Ammonium Nitrogen (NH ₄ -N)	29
Phosphate (P ₂ O ₅)	531
Potash (K ₂ O)	988
Calcium	1,834
Magnesium	280
Sulfur	683
Sodium	94
Aluminum	124
Iron	150
Manganese	17
Copper	6
Zinc	7

⁽¹⁾pH = 6.6; C:N ratio = 13:1.

⁽²⁾Calculation based on applying one-inch thickness of fresh mushroom compost to one acre of land (one acre = 43,560 ft²), which requires approximately 40 tons per acre using an average bulk density of 575 lbs/yd³. For example, applying 40 tons fresh mushroom compost per acre will supply 531 lbs phosphate per acre.



Spent Mushroom Substrate

Spent mushroom substrate is the soil-like material remaining after a crop of mushrooms. Spent substrate is high in organic matter making it desirable for use as a soil amendment or soil conditioner.

 ARTICLES



Sometimes this material is called spent mushroom compost. This fact sheet briefly explains mushroom growing, so that the reader knows what is in the prepared substrate, and then describes the characteristics and possible uses of the material.

Mushroom Growing

Substrate prepared specifically for growing

mushrooms is a blend of natural products. Common ingredients are wheat straw bedding containing horse manure, hay, corn cobs, cottonseed hulls, poultry manure, brewer's grain, cottonseed meal, cocoa bean hulls and gypsum. Growers may add ground soybeans or seed meal supplements later in the production cycle. On top of the substrate, farmers apply a "casing" layer, which is a mixture of peat moss and ground limestone. The casing material provides support for the growing mushrooms.

Spent mushroom substrate still has some nutrients available for the mushroom; however, it is more economical to replace the substrate and start a new crop. Before removing the spent substrate from the mushroom house, the grower "pasteurizes" it with steam to kill any pests or pathogens that may be present in the substrate and casing. This final pasteurization kills weed seeds, insects, and organisms that may cause mushroom diseases. Users may consider spent substrate clean of weed seeds and insects.

Mushroom growers sometimes apply a registered pesticide during the crop cycle. The local garden center sells most of the same pesticides a mushroom farmer uses. Even if pesticides have been applied, they are generally hard to find for two reasons. Organic matter in the substrate effectively binds pesticides. Also, these compounds decompose rapidly at the high temperatures used for pasteurizing the completed crop. It is safe to assume that the pesticide residue on spent substrate is low. Some farms are strictly "organic" and will not use chemical pesticides. These farms can be identified by contacting your Extension office.

Characteristics of Spent Mushroom Substrate

The typical composition of spent mushroom substrate fresh from a mushroom house will vary slightly. Since raw materials and other cultural practices change, each load of fresh spent substrate has a slightly different element and mineral analysis. Therefore the characteristics shown in Table 1 indicate a range of values for each component. Sometimes, fresh substrate is placed in fields for at least one winter season and then marketed as "weathered" mushroom soil. This aged material has slightly different characteristics because the microbial activity in the field will change the composition and texture. The salt content may change during the aging period. If you have any specific questions concerning characteristics of either fresh or aged spent substrate, please contact your local Extension agent.

Appropriate Uses of Spent Substrate

There are many appropriate uses for spent mushroom substrate. Spent mushroom substrate is excellent to spread on top of newly seeded lawns. The material provides cover against birds eating the seeds and will hold the water in the soil while the seeds germinate. Since some plants and garden vegetables are sensitive to high salt content in soils, avoid using fresh spent substrate around those plants. You may use spent substrate weathered for 6 months or longer in all gardens and with most plants. Obtaining spent substrate in the fall and winter, allowing it to weather, will make it ready to use in a garden the following spring. Spring and summer are the best time to use weathered material as a mulch.

As a soil amendment, spent substrate adds organic matter and structure to the soil. Spent substrate primarily improves soil structure and it does provide a few nutrients. Spent substrate is the choice ingredient by those companies making the potting mixtures sold in supermarkets or garden centers. These companies use spent substrate when they need a material to enhance the structure of a soil.

Average Analysis of Spent Mushroom Substrate

Contents	Units	Avg. Fresh	Weathered 16 mos.
Sodium, Na	% Dry Wt.	0.21 - 0.33	0.06
Potassium, K	% Dry Wt.	1.93 - 2.58	0.43
Magnesium, Mg	% Dry Wt.	0.45 - 0.82	0.88
Calcium, Ca	% Dry Wt.	3.63 - 5.15	6.27
Aluminum, Al	% Dry Wt.	0.17 - 0.28	0.58
Iron, Fe	% Dry Wt.	0.18 - 0.34	0.58
Phosphorus, P	% Dry Wt.	0.45 - 0.69	0.84
Ammonia-N, NH ₄	% Dry Wt.	0.06 - 0.24	0.00
Organic Nitrogen	% Dry Wt.	1.25 - 2.15	2.72
Total Nitrogen	% Dry Wt.	1.42 - 2.05	2.72
Solids	% Dry Wt.	33.07 - 40.26	53.47
Volatile Solids	% Dry Wt.	52.49 - 72.42	54.24
pH	Standard Units	5.8 - 7.7	7.1
N-P-K ratio	PPM Dry Wt.	1.8 - 0.6 - 2.2	2.7 - 0.8 - 0.47
% x 10,000 = PPM			

Prepared by David M. Beyer, Penn State

© 2017 Penn State Extension



SAFETY DATA SHEET (SDS) – Mix & Gro SMS-MH

Section 1 - Identification

Product Name: Mix & Gro SMS-MH
Other means of identification: Spent Mushroom Compost.
Recommended use/restrictions: Soil amendment and conditioner. No restrictions
Manufacture: Monterey Mushroom, Inc. - Morgan Hill
642 Hale Ave
Morgan Hill, CA 95037

Section 2 - Hazard(s) identification

Classification: This product is not considered to be hazardous nor to contain hazardous chemicals based on evaluations made by our company under the OSHA Hazard Communication Standard, 29 CFR 1910.1200.

Signal word: No signal word.

Hazard statements:

- May cause eye irritation.
- May cause irritation of respiratory tract if inhaled.
- May cause irritation from prolonged contact with skin.
- Not known to be toxic if ingested.

Symbols: No hazard symbols.

Precautionary statements: Keep out of reach of children. Use as directed.

Section 3 - Composition/Information on ingredients

Chemical name: None.
Common name: Compost.
Ingredients: Spent Mushroom Compost.
CAS#: Not listed in CAS registry.
Manufacturer/distributor: Mix & Gro, LLC
6504 Sutter Avenue
Carmichael, CA 95608
Emergency contact number: (916) 761-6882

Section 4 - First aid measures

Eye contact: Immediately flush thoroughly with water. If irritation persists after five minutes of flushing, remove contact lenses and repeat flushing thoroughly for five more minutes. If irritation persists, get medical attention immediately.

Skin contact: Rinse irritated area of skin with soap and water. If irritation persists, get medical attention immediately.

Inhalation: If irritation is experienced indoors, move outside to fresh air. Rinse mouth with fresh water. If irritation persists, get medical attention immediately.

Ingestion: Drink a glass full of fresh water. If difficulty breathing or nausea is experienced, get medical attention immediately.

Section 5 - Fire-fighting measures

Extinguishing media: Water, foam, chemical or carbon dioxide.

Specific combustion hazards: None.

Precautions for fire-fighters: None.

Section 6 - Accidental release measures

Personal precautions: Wear protective respiratory and eye equipment.

Emergency procedures: None.

Containment and cleanup: Material is non-toxic. No special methods or materials required.

Section 7 - Handling and storage

Safe handling precautions: Wear protective respiratory equipment and safety glasses.

Safe storage: Store in cool, well-ventilated place. No incompatibilities.

Section 8 - Exposure controls/personal protection

Permissible exposure limits: Not controlled under permissible exposure limit (PEL) of OSHA nor Threshold Limit Value (TLV) of American Conference of Governmental Industrial Hygienists.

Engineering controls: Use in well-ventilated area.

Individual protection measures: Wear a NIOSH/MSHA-approved respirator and safety glasses.

Section 9 - Physical and chemical properties

Physical state:	Solid.	Flammability (gas):	Not applicable.
Color:	Dark brown.	Upper/lower flammability or explosive limits:	Not applicable.
Odor:	Earthy.	Vapor pressure:	Not applicable.
Odor threshold:	Not applicable.	Vapor density:	Not applicable.
pH	Neutral.	Relative density:	Not applicable.
Melting/freezing point:	Not applicable.	Solubility:	Insoluble.
Initial boiling point and boiling range:	Not applicable.	Flash point:	Not applicable.
Auto-ignition temperature:	Not applicable.		
Evaporation rate:	Not applicable.	Decomposition temperature:	Not applicable.
Flammability (solid):	Not applicable.	Viscosity:	Not applicable.

Section 10 - Stability and reactivity

Reactivity:	None.
Chemical stability:	Stable.
Hazardous reactions:	None.
Conditions to avoid:	None.
Hazardous decomposition products:	Not applicable.

Section 11 - Toxicological information

Routes of exposure:	Inhalation, ingestion, skin and eye contact.
Immediate effects:	May cause eye irritation. May cause irritation of respiratory tract if inhaled. May cause irritation from prolonged contact with skin. Not known to be toxic if ingested.
Chronic effects:	No known effects.
Toxicity:	No acute toxicity estimates. Not listed in the National Toxicology Program (NTP) Report on Carcinogens nor found to be a potential carcinogen in the International Agency for Research on Cancer (IARC) Monographs or by OSHA.

Section 12 - Ecological information

Ecotoxicity:	No known effects.
Persistence and degradability:	Biodegradable.
Bioaccumulative potential:	Not applicable. Positive effect on soil.
Mobility in soil:	None. Beneficial to soil.
Other:	No known adverse effects.

Section 13 - Disposal considerations

Waste residues:	No product residues. Packaging only.
Residue handling:	Non-toxic, non-contaminated packaging. No special disposal handling required.

Section 14 - Transport information

UN number:	Not applicable.
UN shipping name:	Not applicable.
Transport hazard:	Not regulated by DOT.
Packing group:	Not applicable.
Environmental hazards:	None known.
Bulk transport:	Not applicable.
Special precautions:	None.

Section 15 - Regulatory information

Safety:	Not regulated, but pesticide and pathogen free.
Health:	Not regulated, but pesticide and pathogen free.

Environmental:

Not regulated, but pesticide and pathogen free

Section 16 - Other information

The information provided in this Safety Data Sheet (SDS) is correct to the best of our knowledge and information available at the time of its publication. Information provided is to be utilized only as guidance for safe handling, use, processing, storage, transportation, disposal and release and is not to be considered a warranty or quality assurance or specification. This SDS relates only to Spent Mushroom Compost and may not be valid when used in combination with any other materials or in any process, unless specified in the text.

This SDS was last updated: 11/4/19